

# Linear Programming: Chapter 7

## Sensitivity and Parametric Analysis

Robert J. Vanderbei

October 7, 2010

Slides last edited on October 5, 2010

Operations Research and Financial Engineering  
Princeton University  
Princeton, NJ 08544  
<http://www.princeton.edu/~rvdb>

# Restarting

Consider an optimal dictionary:

$$\begin{aligned}\zeta &= \zeta^* - z_{\mathcal{N}}^{*T} x_{\mathcal{N}} \\ x_{\mathcal{B}} &= x_{\mathcal{B}}^* - B^{-1} N x_{\mathcal{N}}.\end{aligned}$$

Recall definitions of  $x_{\mathcal{B}}^*$ ,  $z_{\mathcal{N}}^*$ , and  $\zeta^*$ :

$$\begin{aligned}x_{\mathcal{B}}^* &= B^{-1} b \\ z_{\mathcal{N}}^* &= (B^{-1} N)^T c_{\mathcal{B}} - c_{\mathcal{N}} \\ \zeta^* &= c_{\mathcal{B}}^T B^{-1} b.\end{aligned}$$

Now, suppose objective coefficients change from  $c$  to  $\tilde{c}$ .

To adjust current dictionary,

- recompute  $z_{\mathcal{N}}^*$ , and
- recompute  $\zeta^*$ .

Note that  $x_{\mathcal{B}}^*$  remains unchanged. Therefore,

- Adjusted dictionary is *primal feasible*.
- Apply primal simplex method.
- Likely to reach optimality quickly.

Had it been the right-hand sides  $b$  that changed, then

- Adjusted dictionary would be *dual feasible*.
- Could apply dual simplex method.

# Ranging

Given an optimal dictionary:

$$\begin{aligned}\zeta &= \zeta^* - z_{\mathcal{N}}^{*T} x_{\mathcal{N}} \\ x_{\mathcal{B}} &= x_{\mathcal{B}}^* - B^{-1} N x_{\mathcal{N}}.\end{aligned}$$

Question: *If  $c$  were to change to*

$$\tilde{c} = c + \mu \Delta c,$$

*for what range of  $\mu$ 's does the current basis remain optimal?*

Recall that:

$$z_{\mathcal{N}}^* = (B^{-1} N)^T c_{\mathcal{B}} - c_{\mathcal{N}}$$

Therefore, dual variables change as follows by  $\mu \Delta z_{\mathcal{N}}$  where

$$\Delta z_{\mathcal{N}} = (B^{-1} N)^T \Delta c_{\mathcal{B}} - \Delta c_{\mathcal{N}}$$

We want:

$$z_{\mathcal{N}}^* + \mu \Delta z_{\mathcal{N}} \geq 0$$

From familiar ratio tests, we get

$$\left( \min_{j \in \mathcal{N}} -\frac{\Delta z_j}{z_j^*} \right)^{-1} \leq \mu \leq \left( \max_{j \in \mathcal{N}} -\frac{\Delta z_j}{z_j^*} \right)^{-1}.$$

Comments:

- A similar analysis works for changes to the right-hand side.
- An example is worked out in the text.

# Ranging with the Pivot Tool.

An initial dictionary:

obj =	0.0	+	2.0	x1	+	1.0	x2	-	-
			-1.0	x1	+	-1.0	x2	-	-
w1 =	-1.0	+	1.0	-	-1.0	x1	-	-1.0	x2
w2 =	2.0	+	1.0	-	-1.0	x1	-	1.0	x2
w3 =	3.0	+	1.0	-	0.0	x1	-	1.0	x2
w4 =	5.0	+	1.0	-	1.0	x1	-	1.0	x2
w5 =	3.0	+	1.0	-	1.0	x1	-	0.0	x2
w6 =	2.0	+	1.0	-	1.0	x1	-	-1.0	x2

The optimal dictionary:

obj =	8.0	+	-1.0	w4	+	-1.0	w5	-	-
			1.0	w4	+	0.0	w5	-	-
w1 =	4.0	+	2.0	-	1.0	w4	-	0.0	w5
w2 =	3.0	+	2.0	-	-1.0	w4	-	2.0	w5
w3 =	1.0	+	1.0	-	-1.0	w4	-	1.0	w5
w6 =	1.0	+	0.0	-	1.0	w4	-	-2.0	w5
x2 =	2.0	+	0.0	-	1.0	w4	-	-1.0	w5
x1 =	3.0	+	1.0	-	0.0	w4	-	1.0	w5

Question: *If the coefficient on  $x_2$  in original problem were changed to  $1 + \mu$  (and everything remains unchanged), for what range of  $\mu$ 's does the current basis remain optimal?*

# Ranging with the Pivot Tool–Continued.

Set artificial rhs column to zeros.

Set artificial objective row to “ $x_2$ ”:

<b>obj</b> =	8.0	+		-	+	-1.0	w4	+	-1.0	w5
				+	+	-1.0	w4	+	1.0	w5
<b>w1</b> =	4.0	+	0.0	-	-	1.0	w4	-	0.0	w5
<b>w2</b> =	3.0	+	0.0	-	-	-1.0	w4	-	2.0	w5
<b>w3</b> =	1.0	+	0.0	-	-	-1.0	w4	-	1.0	w5
<b>w6</b> =	1.0	+	0.0	-	-	1.0	w4	-	-2.0	w5
<b>x2</b> =	2.0	+	0.0	-	-	1.0	w4	-	-1.0	w5
<b>x1</b> =	3.0	+	0.0	-	-	0.0	w4	-	1.0	w5

-1.0     $\Leftarrow \mu \Leftarrow$     1.0

The range of  $\mu$  values is shown at the bottom of the pivot tool.



# Perturb

Introduce a parameter  $\mu$  and perturb:

$$\begin{array}{r}
 \zeta = \qquad \qquad \qquad -3x_1 + 11x_2 + 2x_3 \\
 \qquad \qquad \qquad \qquad \qquad -\mu x_1 - \mu x_2 - \mu x_3 \\
 \hline
 w_1 = 5 + \mu + x_1 - 3x_2 \\
 w_2 = 4 + \mu - 3x_1 - 3x_2 \\
 w_3 = 6 + \mu \qquad \qquad - 3x_2 - 2x_3 \\
 w_4 = -4 + \mu + 3x_1 \qquad \qquad + 5x_3
 \end{array}$$

For  $\mu$  large, dictionary is **optimal**.

Question: For which  $\mu$  values is dictionary optimal? Answer:

$$\begin{array}{r}
 -3 - \mu \leq 0 \\
 11 - \mu \leq 0 \quad * \\
 2 - \mu \leq 0 \quad * \\
 \hline
 5 + \mu \geq 0 \\
 4 + \mu \geq 0 \\
 6 + \mu \geq 0 \\
 -4 + \mu \geq 0 \quad *
 \end{array}$$

Note: only those marked with (\*) give inequalities that omit  $\mu = 0$ .

Tightest:

$$\mu \geq 11$$

Achieved by: objective row perturbation on  $x_2$ .

Let  $x_2$  **enter**.



# Second Pivot

Using the *advanced* pivot tool, the current dictionary is:

obj	=	14.6667	+	-14.0	x1	+	-3.6667	w2	+	2.0	x3
				0.0	x1	+	0.3333	w2	+	-1.0	x3
w1	=	1.0	+	0.0	x1	-	-1.0	w2	-	0.0	x3
x2	=	1.3333	+	0.3333	x1	-	0.3333	w2	-	0.0	x3
w3	=	2.0	+	0.0	x1	-	-1.0	w2	-	2.0	x3
w4	=	-4.0	+	1.0	x1	-	0.0	w2	-	-5.0	x3

Note: the parameter  $\mu$  is not shown. **But it is there!**

Question: For which  $\mu$  values is dictionary optimal? Answer:

$$\begin{array}{rcl}
 -14 & & \leq 0 \\
 -3.67 + 0.33\mu & & \leq 0 \\
 2 - \mu & \leq & 0 \quad * \\
 \hline
 1 & & \geq 0 \\
 1.33 + 0.33\mu & & \geq 0 \\
 2 & & \geq 0 \\
 -4 + \mu & \geq & 0 \quad *
 \end{array}$$

Tightest lower bound:

$$\mu \geq 4$$

Achieved by: constraint containing basic variable  $w_4$ . Let  $w_4$  **leave**.

# Second Pivot–Continued

Who shall enter?

Recall the current dictionary:

obj	=	14.6667	+	-14.0	x1	+	-3.6667	w2	+	2.0	x3	
				0.0	x1	+	0.3333	w2	+	-1.0	x3	
w1	=	1.0	+	0.0	-4.0	x1	-	-1.0	w2	-	0.0	x3
x2	=	1.3333	+	0.3333	1.0	x1	-	0.3333	w2	-	0.0	x3
w3	=	2.0	+	0.0	-3.0	x1	-	-1.0	w2	-	2.0	x3
w4	=	-4.0	+	1.0	-3.0	x1	-	0.0	w2	-	-5.0	x3

Do *dual-type* ratio test using current lowest  $\mu$  value, i.e.  $\mu = 4$ :

$$\begin{aligned} 14 + 0 * 4 - 3y_4 &\geq 0 \\ 3.67 - 0.33 * 4 &\geq 0 \\ -2 + 1 * 4 - 5y_4 &\geq 0 \end{aligned}$$

Tightest:

$$-2 + 1 * 4 - 5y_4 \geq 0.$$

Achieved by: objective term containing nonbasic variable  $x_3$ .

Let  $x_3$  **enter**.

# Third Pivot

The current dictionary is:

obj	=	16.2667	+	-15.2	x1	+	-3.6667	w2	+	0.4	w4			
				0.6	x1	+	0.3333	w2	+	-0.2	w4			
w1	=	1.0	+	0.0		-	-4.0	x1	-	-1.0	w2	-	0.0	w4
x2	=	1.3333	+	0.3333		-	1.0	x1	-	0.3333	w2	-	0.0	w4
w3	=	0.4	+	0.4		-	-4.2	x1	-	-1.0	w2	-	0.4	w4
x3	=	0.8	+	-0.2		-	0.6	x1	-	0.0	w2	-	-0.2	w4

Question: For which  $\mu$  values is dictionary optimal? Answer:

$$\begin{array}{r}
 -15.2 + 0.6\mu \leq 0 \\
 -3.67 + 0.33\mu \leq 0 \\
 0.4 - 0.2\mu \leq 0 \quad * \\
 \hline
 1 \geq 0 \\
 1.33 + 0.33\mu \geq 0 \\
 0.4 + 0.4\mu \geq 0 \\
 0.8 - 0.2\mu \geq 0
 \end{array}$$

Tightest lower bound:

$$\mu \geq 2$$

Achieved by: objective term containing nonbasic variable  $w_4$ .

Let  $w_4$  **enter**.

# Third Pivot–Continued

Who shall leave?

Recall the current dictionary:

obj	=	16.2667		+	-15.2	x1	+	-3.6667	w2	+	0.4	w4
				+	0.6	x1	+	0.3333	w2	+	-0.2	w4
w1	=	1.0	+	0.0	-4.0	x1	-	-1.0	w2	-	0.0	w4
x2	=	1.3333	+	0.3333	1.0	x1	-	0.3333	w2	-	0.0	w4
w3	=	0.4	+	0.4	-4.2	x1	-	-1.0	w2	-	0.4	w4
x3	=	0.8	+	-0.2	0.6	x1	-	0.0	w2	-	-0.2	w4

Do *primal-type* ratio test using current lowest  $\mu$  value, i.e.  $\mu = 2$ :

$$\begin{aligned}
 1 + 0 * 2 &\geq 0 \\
 1.33 + 0.33 * 2 &\geq 0 \\
 0.4 + 0.4 * 2 - 0.4w_4 &\geq 0 \\
 0.8 - 0.2 * 2 + 0.2w_4 &\geq 0
 \end{aligned}$$

Tightest:

$$0.4 + 0.4 * 2 - 0.4w_4 \geq 0$$

Achieved by: constraint containing basic variable  $w_3$ .

Let  $w_3$  leave.

# Fourth Pivot

The current dictionary is:

obj	=	16.6667		+	-11.0	x1	+	-2.6667	w2	+	-1.0	w3	
				+	-1.5	x1	+	-0.1667	w2	+	0.5	w3	
w1	=	1.0	+	0.0	-	-4.0	x1	-	-1.0	w2	-	0.0	w3
x2	=	1.3333	+	0.3333	-	1.0	x1	-	0.3333	w2	-	0.0	w3
w4	=	1.0	+	1.0	-	-10.5	x1	-	-2.5	w2	-	2.5	w3
x3	=	1.0	+	0.0	-	-1.5	x1	-	-0.5	w2	-	0.5	w3

It's **optimal!**

Also, the range of  $\mu$  values includes  $\mu = 0$ :

$$\begin{array}{rcl}
 -11 & - & 1.5\mu \leq 0 \\
 -2.67 & - & 0.167\mu \leq 0 \\
 -1 & + & 0.5\mu \leq 0 \\
 \hline
 1 & & \geq 0 \\
 1.33 & + & 0.33\mu \geq 0 \\
 1 & + & 1\mu \geq 0 \\
 1 & & \geq 0
 \end{array}$$

That is,

$$-1 \leq \mu \leq 2$$

Range of  $\mu$  values is shown at bottom of pivot tool. Invalid ranges are highlighted in yellow.